§ 610.15

identity test shall be specific for each product in a manner that will adequately identify it as the product designated on final container and package labels and circulars, and distinguish it from any other product being processed in the same laboratory. Identity may be established either through the physical or chemical characteristics of the product, inspection by macroscopic or microscopic methods, specific cultural tests, or in vitro or in vivo immunological tests.

§610.15 Constituent materials.

- (a) Ingredients, preservatives, diluents, adjuvants. All ingredients used in a licensed product, and any diluent provided as an aid in the administration of the product, shall meet generally accepted standards of purity and quality. Any preservative used shall be sufficiently nontoxic so that the amount present in the recommended dose of the product will not be toxic to the recipient, and in the combination used it shall not denature the specific substances in the product to result in a decrease below the minimum acceptable potency within the dating period when stored at the recommended temperature. Products in multiple-dose containers shall contain a preservative, except that a preservative need not be added to Yellow Fever Vaccine; Poliovirus Vaccine Live Oral; viral vaccines labeled for use with the jet injector; dried vaccines when the accompanying diluent contains a preservative; or to an Allergenic Product in 50 percent or more volume in volume (v/v) glycerin. An adjuvant shall not be introduced into a product unless there is satisfactory evidence that it does not affect adversely the safety or potency of the product. The amount of aluminum in the recommended individual dose of a biological product shall not exceed:
- 0.85 milligrams if determined by assay;
- (2) 1.14 milligrams if determined by calculation on the basis of the amount of aluminum compound added; or
- (3) 1.25 milligrams determined by assay provided that data demonstrating that the amount of aluminum used is safe and necessary to produce the intended effect are submitted to and approved by the Direc-

tor, Center for Biologics Evaluation and Research.

- (b) Extraneous protein; cell culture produced vaccines. Extraneous protein known to be capable of producing allergenic effects in human subjects shall not be added to a final virus medium of cell culture produced vaccines intended for injection. If serum is used at any stage, its calculated concentration in the final medium shall not exceed 1:1 000 000
- (c) *Antibiotics*. A minimum concentration of antibiotics, other than penicillin, may be added to the production substrate of viral vaccines.

[38 FR 32056, Nov. 20, 1973, as amended at 46 FR 51903, Oct. 23, 1981; 48 FR 13025, Mar. 29, 1983; 48 FR 37023, Aug. 16, 1983; 49 FR 23834, June 8, 1984; 50 FR 4134, Jan. 29, 1985; 51 FR 15607, Apr. 25, 1986; 55 FR 11013, Mar. 26, 1990]

§610.16 Total solids in serums.

Except as otherwise provided by regulation, no liquid serum or antitoxin shall contain more than 20 percent total solids.

§610.17 Permissible combinations.

Licensed products may not be combined with other licensed products either therapeutic, prophylactic or diagnostic, except as a license is obtained for the combined product. Licensed products may not be combined with nonlicensable therapeutic, prophylactic, or diagnostic substances except as a license is obtained for such combination.

§610.18 Cultures.

- (a) Storage and maintenance. Cultures used in the manufacture of products shall be stored in a secure and orderly manner, at a temperature and by a method that will retain the initial characteristics of the organisms and insure freedom from contamination and deterioration.
- (b) Identity and verification. Each culture shall be clearly identified as to source strain. A complete identification of the strain shall be made for each new stock culture preparation. Primary and subsequent seed lots shall be identified by lot number and date of preparation. Periodic tests shall be performed as often as necessary to verify

the integrity of the strain characteristics and freedom from extraneous organisms. Results of all periodic tests for verification of cultures and determination of freedom from extraneous organisms shall be recorded and retained.

- (c) Cell lines used for manufacturing biological products—(1) General requirements. Cell lines used for manufacturing biological products shall be:
 - (i) Identified by history;
- (ii) Described with respect to cytogenetic characteristics and tumorigenicity;
- (iii) Characterized with respect to in vitro growth characteristics and life potential; and
- (iv) Tested for the presence of detectable microbial agents.
- (2) *Tests.* Tests that are necessary to assure the safety, purity, and potency of a product may be required by the Director, Center for Biologics Evaluation and Research.
- (3) Applicability. This paragraph applies to diploid and nondiploid cell lines. Primary cell cultures that are not subcultivated and primary cell cultures that are subsequently subcultivated for only a very limited number of population doublings are not subject to the provisions of this paragraph (c).
- (d) *Records*. The records appropriate for cultures under this section shall be prepared and maintained as required by the applicable provisions of §§ 211.188 and 211.194 of this chapter.

(Approved by the Office of Management and Budget under control number 0910–0139) [38 FR 32056, Nov. 20, 1973, as amended at 51 FR 44453, Dec. 10, 1986; 55 FR 11013, Mar. 26, 1990]

§ 610.19 Status of specific products; Group A streptococcus.

The presence of Group A streptococcus organisms and derivatives of Group A streptococcus in Bacterial Vaccines and Bacterial Antigens with "No U.S. Standard of Potency" may induce dangerous tissue reactions in humans. Available data demonstrate that they are unsafe as ingredients in products for human use. Group A streptococcus organisms and derivatives of Group A streptococcus are prohibited from Bacterial Vaccines and Bacterial Antigens with "No U.S. Standard of Potency." Any Bacterial Vaccine or

Bacterial Antigen with "No U.S. Standard of Potency" containing Group A streptococcus organisms or derivatives of Group A streptococcus in interstate commerce is in violation of section 351 of the Public Health Service Act (42 U.S.C. 262).

[44 FR 1549, Jan. 5, 1979]

Subpart C—Standard Preparations and Limits of Potency

§610.20 Standard preparations.

Standard preparations made available by the Center for Biologics Evaluation and Research shall be applied in testing, as follows:

(a) *Potency standards*. Potency standards shall be applied in testing for potency all forms of the following:

ANTIBODIES

Botulism Antitoxin, Type A. Botulism Antitoxin, Type B. Botulism Antitoxin, Type B. Botulism Antitoxin, Type E. Diphtheria Antitoxin.
Histolyticus Antitoxin.
Oedematiens Antitoxin.
Perfringens Antitoxin.
Antipertussis Serum.
Antirabies Serum.
Sordellii Antitoxin.
Staphylococcus Antitoxin.
Tetanus Antitoxin.
Vibrion Septique Antitoxin.

ANTIGENS

Cholera Vaccine, Ogawa serotype.
Diphtheria Toxin for Schick Test.
Pertussis Vaccine.
Tuberculin, Old.
Tuberculin, Purified Protein Derivative.
Typhoid Vaccine.

Cholera Vaccine, Inaba serotype.

BLOOD DERIVATIVE

Thrombin.

(b) Opacity standard. The U.S. Opacity Standard shall be applied in estimating the bacterial concentration of all bacterial vaccines. The assigned value of the standard when observed visually is 10 units. The assigned value of the standard when observed with a photometer is (1) 10 units when the wavelength of the filter is 530 millimicrons, (2) 10.6 units when the wavelength of the filter is 650 millimicrons,